

The analysis of reactions $\pi N \rightarrow \text{two mesons} + N$ within reggeon exchanges.

2. Basic formulas for fit.

V.V. Anisovich and A.V. Sarantsev

Petersburg Nuclear Physics Institute, Gatchina, 188300, Russia

March 3, 2009

Abstract

We present technical aspects of the fitting procedure given in the paper V.V. Anisovich and A.V. Sarantsev *The analysis of reactions $\pi N \rightarrow \text{two mesons} + N$ within reggeon exchanges. 1. Fit and results.*

PACS numbers: 11.25.Hf, 123.1K

1 Introduction

The novel point of the analysis given in the paper [1] is a direct use of reggeon exchange technique for the description of the reactions $\pi N \rightarrow \text{two mesons} + N$ at large energies of the initial pion. This approach allows us to describe simultaneously distributions over M (invariant mass of two mesons) and t (momentum transfer squared to nucleons). Making use of this technique, the following resonances (as well as corresponding bare states), produced in the $\pi N \rightarrow \pi^0 \pi^0 N$ reaction were studied: $f_0(980)$, $f_0(1300)$, $f_0(1200 - 1600)$, $f_0(1500)$, $f_0(1750)$, $f_2(1270)$, $f_2(1525)$, $f_2(1565)$, $f_2(2020)$, $f_4(2025)$. Adding the data on the processes $p\bar{p}(\text{at rest, from liquid H}_2) \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \pi^0 \eta$, $\pi^0 \eta \eta$ and $p\bar{p}(\text{at rest, from gaseous H}_2) \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \pi^0 \eta$, $\pi^0 \eta \eta$, we performed simultaneous K -matrix fit of two-meson spectra in all these reactions. In [1], the results of the combined fit to the above-listed isoscalar f_J -states and isovector ones, $a_0(980)$, $a_0(1475)$, $a_2(1320)$, were presented.

Here we discuss technical aspects of the fitting procedure used in [1]. To give the guide to a reader through the states, which are discussed in [1], the assignment of mesons to Regge trajectories as well as to $q\bar{q}$ nonets is presented following Ref. [2].

The paper is organized as follows.

In Section 2, we present the necessary elements of the reggeon exchange technique in the two-meson production reactions. In Section 3, we discuss the status of the Regge trajectories on (J, M^2) -plane. In Section 4, we give the connection between the dispersion integral equation for a three-body system and the K -matrix approach. We demonstrate what type of assumptions is needed to transform the three-body dispersion relation amplitudes into those of the K -matrix approach. The assignment of mesons to $q\bar{q}$ nonets is given in Section 5.

2 Elements of the reggeon exchange technique in the two-meson production reactions

Here we present the details for the partial wave analysis of two-meson system produced in the high energy πN interaction when the two-meson production occurs due to reggeon exchanges. The reggeon exchange approach is a good tool for studying hadron binary reactions and processes with diffractive production of hadrons at high energies (see [3], Chapters 2 and 6). Interference effects in the amplitudes of the type $\pi N \rightarrow \text{two mesons} + N$ provide valuable information on the contributions of resonances with different quantum numbers.

2.0.1 Kinematics for reggeon exchange amplitudes

For illustration, we consider the reaction $\pi^- p \rightarrow \pi\pi + n$ in the c.m. system of the reaction and present the momenta of the incoming and outgoing particles (below we use the notation for four-vectors: $x = (x_0, \vec{x}_\top, x_z)$).

For the incoming particles we have:

$$\begin{aligned} \text{pion momentum} & : & p_1 &= (p_z + \frac{m_\pi^2}{2p_z}, 0, p_z), \\ \text{proton momentum} & : & p_2 &= (p_z + \frac{m_N^2}{2p_z}, 0, -p_z), \\ \text{total energy squared} & : & s_{\pi N} &= (p_1 + p_2)^2. \end{aligned} \tag{1}$$

Here we have performed an expansion over the large momentum p_z . Analogously, we write for the outgoing particles:

$$\begin{aligned} \text{meson momenta } (i = 1, 2) & : & k_i &= (k_{iz} - \frac{m_i^2 + k_{i\top}^2}{2k_{iz}}, \vec{k}_{i\top}, k_{iz}), \\ \text{total momentum of mesons} & : & P &= k_1 + k_2 = p_1 - q = (p_z + \frac{s + m_\pi^2 + 2q_\top^2}{4p_z}, \vec{q}_\top, p_z - \frac{s - m_\pi^2}{4p_z}), \\ \text{proton momentum} & : & k_3 &= p_2 + q = (p_z - \frac{s - m_\pi^2 + 2q_\top^2}{4p_z}, -\vec{q}_\top, -p_z + \frac{s - m_\pi^2}{4p_z}), \\ \text{energy squared of mesons} & : & s &= P^2 = (k_1 + k_2)^2. \end{aligned} \tag{2}$$

The relative momenta of mesons in the initial and final states read:

$$p = \frac{1}{2}(p_1 + q), \quad k = \frac{1}{2}(k_1 - k_2). \quad (4)$$

The momentum squared transferred to the nucleon is comparatively small:
 $t \equiv q^2 \sim m_N^2 \ll s_{\pi N}$ where

$$q = \left(-\frac{s + m_\pi^2 + 2q_\top^2}{4p_z}, -\vec{q}_\top, \frac{s - m_\pi^2}{4p_z} \right). \quad (5)$$

Neglecting $O(1/p_z^2)$ -terms, one has $q \simeq (0, -\vec{q}_\top, 0)$ and $q^2 \simeq -q_\top^2$.

2.0.2 Angular momentum operators for two-meson systems

As in Ref. [1], we use angular momentum operators $X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp)$, $Z_{\mu_1 \dots \mu_L}^\alpha(k^\perp)$ and projection operator $O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(\perp P)$. Recall their definition.

The operators are constructed from the relative momenta $k_\mu^\perp$ and tensor $g_{\mu\nu}^\perp$. Both of them are orthogonal to the total momentum of the system:

$$k_\mu^\perp = \frac{1}{2}g_{\mu\nu}^\perp(k_1 - k_2)_\nu = k_{1\nu}g_{\nu\mu}^{\perp P} = -k_{2\nu}g_{\nu\mu}^{\perp P}, \quad g_{\mu\nu}^\perp = g_{\mu\nu} - \frac{P_\mu P_\nu}{s}. \quad (6)$$

The operator for $L = 0$ is a scalar (we write $X^{(0)}(k^\perp) = 1$), and the operator for $L = 1$ is a vector, $X_\mu^{(1)} = k_\mu^\perp$. The operators $X_{\mu_1 \dots \mu_L}^{(L)}$ for $L \geq 1$ can be written in the form of a recurrency relation:

$$\begin{aligned} X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) &= k_\alpha^\perp Z_{\mu_1 \dots \mu_L}^\alpha(k^\perp) \equiv k_\alpha^\perp Z_{\mu_1 \dots \mu_L, \alpha}(k^\perp), \\ Z_{\mu_1 \dots \mu_L}^\alpha(k^\perp) &\equiv Z_{\mu_1 \dots \mu_L, \alpha}(k^\perp) = \frac{2L-1}{L^2} \left(\sum_{i=1}^L X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L}^{(L-1)}(k^\perp) g_{\mu_i \alpha}^\perp \right. \\ &\quad \left. - \frac{2}{2L-1} \sum_{\substack{i,j=1 \\ i < j}}^L g_{\mu_i \mu_j}^\perp X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L}^{(L-1)}(k^\perp) \right). \end{aligned} \quad (7)$$

We have a convolution equality $X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) k_{\mu_L}^\perp = k_\perp^2 X_{\mu_1 \dots \mu_{L-1}}^{(L-1)}(k^\perp)$, with $k_\perp^2 \equiv k_\mu^\perp k_\mu^\perp$, and the tracelessness property of $X_{\mu\mu\mu_1 \dots \mu_L}^{(L)} = 0$. On this basis, one can write down the orthogonality-normalization condition for orbital angular operators:

$$\int \frac{d\Omega}{4\pi} X_{\mu_1 \dots \mu'_L \dots \mu_L}^{(L)}(k^\perp) X_{\mu_1 \dots \mu'_L}^{(L')}(k^\perp) = \delta_{LL'} \alpha_L k_\perp^{2L}, \quad \alpha_L = \prod_{l=1}^L \frac{2l-1}{l}, \quad (8)$$

where integration is performed over spherical variables: $\int d\Omega/(4\pi) = 1$.

Iterating equation (7), one obtains the following expression for the operator $X_{\mu_1 \dots \mu_L}^{(L)}$:

$$\begin{aligned} X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) &= \alpha_L \left[k_{\mu_1}^\perp k_{\mu_2}^\perp k_{\mu_3}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp \right. \\ &\quad - \frac{k_\perp^2}{2L-1} \left(g_{\mu_1 \mu_2}^\perp k_{\mu_3}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp + g_{\mu_1 \mu_3}^\perp k_{\mu_2}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp + \dots \right) \\ &\quad \left. + \frac{k_\perp^4}{(2L-1)(2L-3)} \left(g_{\mu_1 \mu_2}^\perp g_{\mu_3 \mu_4}^\perp k_{\mu_5}^\perp k_{\mu_6}^\perp \dots k_{\mu_L}^\perp + g_{\mu_1 \mu_2}^\perp g_{\mu_3 \mu_5}^\perp k_{\mu_4}^\perp k_{\mu_6}^\perp \dots k_{\mu_L}^\perp + \dots \right) + \dots \right]. \end{aligned} \quad (9)$$

For the projection operators, one has:

$$\begin{aligned} O &= 1, \quad O_\nu^\mu(\perp P) = g_{\mu\nu}^\perp, \\ O_{\nu_1 \nu_2}^{\mu_1 \mu_2}(\perp P) &= \frac{1}{2} \left(g_{\mu_1 \nu_1}^\perp g_{\mu_2 \nu_2}^\perp + g_{\mu_1 \nu_2}^\perp g_{\mu_2 \nu_1}^\perp - \frac{2}{3} g_{\mu_1 \mu_2}^\perp g_{\nu_1 \nu_2}^\perp \right). \end{aligned} \quad (10)$$

For higher states, the operator can be calculated using the recurrent expression:

$$\begin{aligned} O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(\perp P) &= \frac{1}{L^2} \left(\sum_{i,j=1}^L g_{\mu_i \nu_j}^\perp O_{\nu_1 \dots \nu_{j-1} \nu_{j+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L}(\perp P) \right. \\ &\quad \left. - \frac{4}{(2L-1)(2L-3)} \times \sum_{\substack{i < j \\ k < m}}^L g_{\mu_i \mu_j}^\perp g_{\nu_k \nu_m}^\perp O_{\nu_1 \dots \nu_{k-1} \nu_{k+1} \dots \nu_{m-1} \nu_{m+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L}(\perp P) \right). \end{aligned} \quad (11)$$

The projection operators obey the relations:

$$\begin{aligned} O_{\nu_1 \dots \nu_J}^{\mu_1 \dots \mu_J}(\perp P) X_{\nu_1 \dots \nu_J}^{(J)}(k^\perp) &= X_{\mu_1 \dots \mu_J}^{(J)}(k^\perp), \\ O_{\nu_1 \dots \nu_J}^{\mu_1 \dots \mu_J}(\perp P) k_{\nu_1} k_{\nu_2} \dots k_{\nu_J} &= \frac{1}{\alpha_J} X_{\mu_1 \dots \mu_J}^{(J)}(k^\perp). \end{aligned} \quad (12)$$

Hence, the product of the two $X^J(k_\perp)$ operators results in the Legendre polynomials as follows:

$$X_{\mu_1 \dots \mu_J}^{(J)}(k_\perp)(p^\perp)(-1)^J O_{\nu_1 \dots \nu_J}^{\mu_1 \dots \mu_J}(\perp P) X_{\nu_1 \dots \nu_J}^{(J)}(k^\perp) = \alpha_J (\sqrt{-p_\perp^2} \sqrt{-k_\perp^2})^J P_J(z), \quad (13)$$

where $z \equiv (-p^\perp k^\perp) / (\sqrt{-p_\perp^2} \sqrt{-k_\perp^2})$.

2.1 Reggeized pion exchanges

To be definite, we present here formulas which lead to differential cross-section moment expansion in processes initiated by reggeized pions situated on the trajectories $R(\pi_j)$. Recall that index j labels different trajectories, which are leading and daughter ones, as well as trajectories generated by a reggeization of the t -channel states with $J > 0$ (in multiparticle processes, like $\pi N \rightarrow \text{two mesons} + N$, the vertices $\pi R(\pi_j)$ are different for states with $J = 0$ and $J > 0$, so it is convenient to separate their contributions).

2.1.1 Production of two mesons at small $|t|$ – the hypothesis of dominant pion exchange

Under this hypothesis, the amplitude for the $\pi\pi$ production block is written as follows:

$$A(\pi R(\pi_j) \rightarrow \pi\pi) \rightarrow A(\pi\pi \rightarrow \pi\pi) = 16\pi \sum_J A_{\pi\pi \rightarrow \pi\pi}^J(s) (2J+1) N_J^0 Y_J^0(z, \varphi), \quad (14)$$

$$Y_J^m(z, \varphi) = \frac{1}{N_J^m} P_J^m(z) e^{im\varphi}, \quad N_J^m = \sqrt{\frac{4\pi}{2J+1} \frac{(J+m)!}{(J-m)!}}.$$

Below we consider the resonance decay in the set of channels with two pseudoscalar mesons in the final states, that means $\pi\pi \rightarrow c = \pi\pi, K\bar{K}, \eta\eta, \dots$. Generalizing consideration, we give the same freedom for initial state. Therefore, we denote the initial and final ($I = 0$)-channel states as a, c with $a, c = \pi\pi, K\bar{K}, \eta\eta$, and so on. Then the transition amplitudes are denoted as

$$X_{\mu_1 \dots \mu_J}^{(J)}(p^\perp) A_{a \rightarrow c}^J(s) (-1)^J O_{\nu_1 \dots \nu_J}^{\mu_1 \dots \mu_J}(\perp P) X_{\nu_1 \dots \nu_J}^{(J)}(k_c^\perp) \xi_J, \quad \xi_J = \frac{16\pi(2J+1)}{\alpha_J}, \quad (15)$$

where $k_{c\mu}^\perp = \frac{1}{2}(k_{c1\nu}^\perp - k_{c2\nu}^\perp)g_{\nu\mu}^\perp$.

The unitarity condition for the transition amplitudes reads:

$$\text{Im} A_{a \rightarrow c}^J(s) = \sum_b \frac{2\sqrt{-k_{b\perp}^2}}{\sqrt{s}} A_{a \rightarrow b}^J(s) A_{b \rightarrow c}^{J*}(s) (-k_{c\perp}^2)^J, \quad (16)$$

where index b refers to intermediate states ($b = \pi\pi, K\bar{K}, \eta\eta, \dots$). The unitarity condition is fulfilled by using for $A_{a \rightarrow c}^J(s)$ the K -matrix form:

$$A_{a \rightarrow c}^J(s) = \sum_b \hat{K}_{ab}^J \left(\frac{I}{I - i\hat{\rho}^J(s)\hat{K}^J} \right)_{bc}, \quad (17)$$

where $\hat{\rho}$ is a diagonal matrix with elements $\rho_{bb}^J(s) = 2\sqrt{-k_{b\perp}^2} (-k_{b\perp}^2)^J / \sqrt{s}$.

We parametrize the elements of the K -matrix in the following form:

$$\begin{aligned} K_{ab}^J &= \sum_n \frac{1}{B_J(-k_{a\perp}^2, r_n)} \left(\frac{g_a^{n(J)} g_b^{n(J)}}{M_n^2 - s} \right) \frac{1}{B_J(-k_{b\perp}^2, r_n)} \\ &+ \frac{f_{ab}^{(J)}}{B_J(-k_{a\perp}^2, r_0) B_J(-k_{b\perp}^2, r_0)}. \end{aligned} \quad (18)$$

Index n refers to a set of resonances, the resonance couplings g_c^n are constants, and f_{ac} is a non-resonance term. The form factors $B_J(-k_\perp^2, r)$ are introduced to compensate the divergence of the relative momentum factor at large energies. Such form factors are known as the Blatt–Weisskopf factors depending on the radius of the state r_n . For non-resonance transition, the radius r_0 is taken to be much larger than that for resonance contributions.

2.1.2 Calculation routine for the reggeized pion exchanges

In case of the two-meson production, the initial-state K -matrix element is called the P -vector: $K_{\pi R(\pi_j),b}^J \equiv P_{\pi R(\pi_j),b}^J$ (recall that index j labels the leading and daughter trajectories). We write for the two-meson production amplitude initiated by the pion reggeon exchange the following representation:

$$A_{\pi R(\pi_j),c}^J(s) = \sum_b P_{\pi R(\pi_j),b}^J \left(\frac{I}{I - i\hat{\rho}^J(s)\hat{K}^J} \right)_{bc}. \quad (19)$$

The P -vector is parametrized in the form similar to eq. (18):

$$\begin{aligned} P_{\pi R(\pi_j),c}^J &= \sum_n \frac{1}{B_J(-p_\perp^2, r_n)} \left(\frac{G_{\pi R(\pi_j)}^{n(J)}(q^2) g_c^{n(J)}}{M_n^2 - s} \right) \frac{1}{B_J(-k_{c\perp}^2, r_n)} \\ &+ \frac{F_{\pi R(\pi_j),c}^{(J)}(q^2)}{B_J(-p_\perp^2, r_0) B_J(-k_{c\perp}^2, r_0)}. \end{aligned} \quad (20)$$

The product of the two amplitudes is equal to:

$$\begin{aligned} A(\pi R(\pi_j) \rightarrow \pi\pi) A^*(\pi R(\pi_k) \rightarrow \pi\pi) &= (16\pi)^2 \sum_J Y_J^0(z, \varphi) \\ &\times \sum_{J_1 J_2} d_{J_1 J_2 J}^{000} A_{\pi R(\pi_j) \rightarrow \pi\pi}^{J_1}(s) A_{\pi R(\pi_k) \rightarrow \pi\pi}^{J_2}(s) (2J_1+1)(2J_2+1) N_{J_1}^0 N_{J_2}^0, \end{aligned} \quad (21)$$

where the coefficients d_{nmk}^{iji+j} are given below. Averaging over the polarizations of the initial nucleons and summing over polarisation of the final ones, we get $\text{Sp}[(\vec{\sigma}\vec{q}_\perp)(\vec{\sigma}\vec{q}_\perp)] \simeq -q^2 = -t$. So, we obtain for the total amplitude squared:

$$\begin{aligned} |A_{\pi p \rightarrow \pi\pi n}^{\text{(pion trajectories)}}|^2 &= \sum_{R(\pi_j) R(\pi_k)} A(\pi R(\pi_j) \rightarrow \pi\pi) A^*(\pi R(\pi_k) \rightarrow \pi\pi) \\ &\times R_{\pi_j}(s_{\pi N}, q^2) R_{\pi_k}^*(s_{\pi N}, q^2) (-t) (g_{pn}^{(\pi)})^2. \end{aligned} \quad (22)$$

The final expression reads:

$$\begin{aligned} N(M, t) \langle Y_J^0 \rangle &= \frac{\rho(s) \sqrt{s}}{\pi |\vec{p}_2|^2 s_{\pi N}} \sum_{R(\pi_j) R(\pi_k)} R_{\pi_j}(s_{\pi N}, q^2) R_{\pi_k}^*(s_{\pi N}, q^2) (-t) (g_{pn}^{(\pi)})^2 \\ &\times \sum_{J_1 J_2} d_{J_1 J_2 J}^{000} A_{\pi\pi_j \rightarrow \pi\pi}^{J_1}(s) A_{\pi R(\pi_k) \rightarrow \pi\pi}^{J_2}(s) (2J_1+1)(2J_2+1) N_{J_1}^0 N_{J_2}^0. \end{aligned} \quad (23)$$

(i) Spherical functions.

Let us present here some relations for the spherical functions used in the calculations:

$$\begin{aligned} Y_l^m(\Theta, \varphi) &= \frac{1}{N_l^m} P_l^m(z) e^{im\varphi}, \quad N_l^m = \sqrt{\frac{4\pi}{2l+1} \frac{(n+m)!}{(l-m)!}}, \\ P_l^m(z) &= (-1)^m (1-z^2)^{\frac{m}{2}} \frac{d^m}{dz^m} P_l(z), \end{aligned} \quad (24)$$

where $z = \cos \Theta$. We have the following convolution rule for two spherical functions:

$$Y_n^i(\Theta, \varphi) Y_m^j(\Theta, \varphi) = \sum_{k=0}^{n+m} d_{n,m,k}^{i,j,i+j} Y_k^{i+j}(\Theta, \varphi), \quad (25)$$

$d_{n,m,k}^{i,j,i+j}$ is defined also in [1], eq. (8).

2.1.3 Calculations related to the expansion of the differential cross section $\pi p \rightarrow \pi\pi + N$ over spherical functions for the reggeised π_2 -exchange

Here we present formulas which refer to the calculation routine related to the reggeised π_2 -exchange.

The convolution of angular momentum operators can be expressed through Legendre polynomials and their derivatives:

$$\begin{aligned} & X_{\alpha\beta\mu_1\ldots\mu_J}^{(J+2)}(p^\perp) (-1)^J O_{\nu_1\ldots\nu_J}^{\mu_1\ldots\mu_J}(\perp P) X_{\nu_1\ldots\nu_J}^{(J)}(k^\perp) \\ &= \frac{2\alpha_J \left(\sqrt{-k_\perp^2}\right)^J \left(\sqrt{-p_\perp^2}\right)^{J+2}}{3(J+1)(J+2)} \\ &\times \left(X_{\mu\nu}^{(2)}(p^\perp) \frac{P_{J+2}''}{-p_\perp^2} + X_{\mu\nu}^{(2)}(k^\perp) \frac{P_J''}{-k_\perp^2} - \frac{3 P_{J+1}''}{\sqrt{-k_\perp^2} \sqrt{-p_\perp^2}} k_\mu^\perp p_\nu^\perp \right) O_{\mu\nu}^{\alpha\beta}(\perp q), \\ & O_{\chi\tau}^{\alpha\beta}(\perp q) X_{\chi\mu_2\ldots\mu_J}^{(J)}(p^\perp) (-1)^J O_{\nu_1\nu_2\ldots\nu_J}^{\tau\mu_2\ldots\mu_J}(\perp P) X_{\nu_1\ldots\nu_J}^{(J)}(k^\perp) \\ &= \frac{2\alpha_{J-1}}{3J^2} \left(\sqrt{-k_\perp^2}\right)^J \left(\sqrt{-p_\perp^2}\right)^J \\ &\times \left(X_{\mu\nu}^{(2)}(p^\perp) \frac{P_J''}{-p_\perp^2} + X_{\mu\nu}^{(2)}(k^\perp) \frac{P_J''}{-k_\perp^2} - \frac{P_J' + 2z P_J''}{\sqrt{-k_\perp^2} \sqrt{-p_\perp^2}} k_\mu^\perp p_\nu^\perp \right) O_{\mu\nu}^{\alpha\beta}(\perp q), \\ & X_{\mu_3\ldots\mu_J}^{(J-2)}(p^\perp) (-1)^J O_{\nu_1\nu_2\nu_3\ldots\nu_J}^{\alpha\beta\mu_3\ldots\mu_J}(\perp P) X_{\nu_1\ldots\nu_J}^{(J)}(k^\perp) \\ &= \frac{2\alpha_{J-2} \left(\sqrt{-k_\perp^2}\right)^J \left(\sqrt{-p_\perp^2}\right)^{J-2}}{3(n-1)n} \\ &\times \left(X_{\mu\nu}^{(2)}(p^\perp) \frac{P_{J-2}''}{-p_\perp^2} + X_{\mu\nu}^{(2)}(k^\perp) \frac{P_J''}{-k_\perp^2} - \frac{3 P_{J-1}''}{\sqrt{-k_\perp^2} \sqrt{-p_\perp^2}} k_\mu^\perp p_\nu^\perp \right) O_{\mu\nu}^{\alpha\beta}(\perp q). \end{aligned} \quad (26)$$

Let us remind that $p_\mu^\perp = p_{1\nu} g_{\nu\mu}^{\perp P}$ and $k_\mu^\perp = k_{1\nu} g_{\nu\mu}^{\perp P}$. Therefore the amplitude $A_{\alpha\beta}(\pi R(\pi_2) \rightarrow \pi\pi)$ can be rewritten as:

$$\begin{aligned} A_{\alpha\beta}(\pi R(\pi_2) \rightarrow \pi\pi) &= \frac{2}{3} \sum_J \left[\frac{X_{\alpha\beta}^{(2)}(p^\perp)}{-p_\perp^2} \left(C_1^{(J)} P_{J+2}'' A_{+2}^{(J)}(s) + \right. \right. \\ &\left. \left. + C_2^{(J)} P_J'' A_0^{(J)}(s) + C_3^{(J)} P_{J-2}'' A_{-2}^{(J)}(s) \right) \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{X_{\alpha\beta}^{(2)}(k^\perp)}{-k_\perp^2} P_J'' \left(C_1^{(J)} A_{+2}^{(J)}(s) + C_2^{(J)} A_0^{(J)}(s) + C_3^{(J)} A_{-2}^{(J)}(s) \right) \\
& - \frac{O_{\mu\nu}^{\alpha\beta} k_\mu^\perp p_\nu^\perp}{\sqrt{-k_\perp^2} \sqrt{-p_\perp^2}} \left(3C_1^{(J)} P_{J+1}'' A_{+2}^{(J)}(s) + C_2^{(J)} (P_J' + 2zP_J'') A_0^{(J)}(s) \right. \\
& \left. + 3C_3^{(J)} P_{J-1}'' A_{-2}^{(J)}(s) \right) \Big], \tag{27}
\end{aligned}$$

where

$$\begin{aligned}
C_1^{(J)} &= \frac{16\pi(2J+1)}{(J+1)(J+2)} \left(\sqrt{-k_\perp^2} \right)^J \left(\sqrt{-p_\perp^2} \right)^{J+2}, \\
C_2^{(J)} &= \frac{16\pi(2J+1)}{J(2J-1)} \left(\sqrt{-k_\perp^2} \right)^J \left(\sqrt{-p_\perp^2} \right)^J, \\
C_3^{(J)} &= \frac{16\pi(2J+1)}{(2J-1)(2J-3)} \left(\sqrt{-k_\perp^2} \right)^J \left(\sqrt{-p_\perp^2} \right)^{J-2}. \tag{28}
\end{aligned}$$

In the amplitude with the $X_{\alpha\beta}^{(2)}(p^\perp)$ structure there is no $m = 1$ component. This amplitude should be taken effectively into account by the π trajectory. The second amplitude has the same angular dependence $P_J''(z)$ and works for resonances with $J \geq 2$. In the first approximation it is reasonable to use the third term only, which has the smallest power of $p_\perp^2$.

The third amplitude has angular dependencies:

$$P_{J+1}''(z), \quad P_J' + 2zP_J'', \quad P_{J-1}''. \tag{29}$$

The first and second angular dependencies are the same for $J = 1, 2$ and differ only at $n \geq 3$, when the third term appears. Therefore, in the first approximation one can use only the second term which has a lower order of $p_\perp^2$ to fit the data. Thus, the $R(\pi_2)$ exchange amplitude can be approximated as:

$$\begin{aligned}
A_{\alpha\beta}(\pi R(\pi_2) \rightarrow \pi\pi) &\simeq \frac{2}{3} \sum_J \left[\frac{X_{\alpha\beta}^{(2)}(k^\perp)}{-k_\perp^2} P_J'' C_3^{(J)} A_{-2}^{(J)}(s) \right. \\
&\quad \left. - \frac{O_{\mu\nu}^{\alpha\beta} k_\mu^\perp p_\nu^\perp}{\sqrt{-k_\perp^2} \sqrt{-p_\perp^2}} C_2^{(J)} (P_J' + 2zP_J'') A_0^{(J)}(s) \right]. \tag{30}
\end{aligned}$$

The convolution of operators in (30) with $k_{3\alpha}^{\perp q} k_{3\beta}^{\perp q}$ in the GJ system gives:

$$\begin{aligned}
k_{3\alpha}^{\perp q} k_{3\beta}^{\perp q} X_{\alpha\beta}^{(2)}(k_\perp^2) &= |\vec{k}|^2 k_{3z}^{\perp q} \left(k_{3z}^{\perp q} P_2(z) + 3k_{3x} z \cos \varphi \sin \Theta \right), \\
k_{3\alpha}^{\perp q} k_{3\beta}^{\perp q} O_{\mu\nu}^{\alpha\beta} k_\mu^\perp p_\nu^\perp &= \frac{1}{3} |\vec{k}| |\vec{p}| k_{3z}^{\perp q} \left(2k_{3z}^{\perp q} z + 3k_{3x}^{\perp q} \cos \varphi \sin \Theta \right), \tag{31}
\end{aligned}$$

and the total amplitude is equal to:

$$\begin{aligned}
A_{\pi p \rightarrow \pi \pi n}^{R(\pi_2)} &= \frac{1}{s_{\pi N}^2} \sum_J \left(V_1^J A_{-2}^{(J)}(s) - V_2^J A_0^{(J)}(s) \right) R_{\pi_2}(s_{\pi N}, q^2) \\
&\times \left(\varphi_n^+(\vec{\sigma} \vec{p}_\perp) \varphi_p \right) g_{pn}^{(\pi_2)}, \tag{32}
\end{aligned}$$

where

$$\begin{aligned} V_1^{(J)} &= C_3^{(J)} k_{3z}^{\perp q} \left(k_{3z}^{\perp q} P_2(z) + 3k_{3x} z \cos \varphi \sin \Theta \right) P_J'', \\ V_2^{(J)} &= \frac{1}{3} C_2^{(J)} k_{3z}^{\perp q} (P_J' + 2z P_J'') \left(2k_{3z}^{\perp q} z + 3k_{3x}^{\perp q} \cos \varphi \sin \Theta \right). \end{aligned} \quad (33)$$

For $J = 1$ the first vertex is equal to 0; for the next ones we have:

$$\begin{aligned} V_2^{(1)} &= \frac{1}{3} C_2^{(1)} k_{3z}^{\perp q} \left(2k_{3z}^{\perp q} Y_1^0 N_1^0 - 3k_{3x}^{\perp q} \text{Re } Y_1^1 N_1^1 \right), \\ V_1^{(2)} &= C_3^{(2)} k_{3z}^{\perp q} \left(k_{3z}^{\perp q} Y_2^0 N_2^0 - k_{3x} \text{Re } Y_2^1 N_2^1 \right), \\ V_2^{(2)} &= \frac{1}{3} C_2^{(2)} k_{3z}^{\perp q} \left(12k_{3z}^{\perp q} Y_2^0 N_2^0 + 6k_{3z}^{\perp q} Y_0^0 N_0^0 - 9k_{3x}^{\perp q} \text{Re } Y_2^1 N_2^1 \right), \\ V_1^{(3)} &= C_3^{(3)} k_{3z}^{\perp q} \left(9k_{3z}^{\perp q} Y_3^0 N_3^0 + 18k_{3z}^{\perp q} Y_1^0 N_1^0 - 6k_{3x} \text{Re } Y_3^1 N_3^1 - 9k_{3x} \text{Re } Y_1^1 N_1^1 \right), \\ V_2^{(3)} &= \frac{1}{3} C_2^{(3)} k_{3z}^{\perp q} \left(30k_{3z}^{\perp q} Y_3^0 N_3^0 + 42k_{3z}^{\perp q} Y_1^0 N_1^0 - 15k_{3x}^{\perp q} \text{Re } Y_3^1 N_3^1 \right. \\ &\quad \left. - 18k_{3x}^{\perp q} \text{Re } Y_1^1 N_1^1 \right), \\ V_1^{(4)} &= C_3^{(4)} k_{3z}^{\perp q} \left(18k_{3z}^{\perp q} Y_4^0 N_4^0 + 20k_{3z}^{\perp q} Y_2^0 N_2^0 + 7k_{3z}^{\perp q} Y_0^0 N_0^0 \right. \\ &\quad \left. - 9k_{3x} \text{Re } Y_4^1 N_4^1 - 15k_{3x} \text{Re } Y_2^1 N_2^1 \right), \\ V_2^{(4)} &= \frac{C_2^{(4)}}{3} k_{3z}^{\perp q} \left[k_{3z}^{\perp q} \left(56Y_4^0 N_4^0 + 110Y_2^0 N_2^0 + 34Y_0^0 N_0^0 \right) \right. \\ &\quad \left. - k_{3x}^{\perp q} \left(21 \text{Re } Y_4^1 N_4^1 - 30 \text{Re } Y_2^1 N_2^1 \right) \right]. \end{aligned} \quad (34)$$

In a general form, the expression can be written as:

$$\begin{aligned} V_1^{(J)} &= \sum_{n=0}^J C_3^{(J)} k_{3z}^{\perp q} \left[k_{3z}^{\perp q} Y_n^0 R_n^0 (P_2 P_J'') + 3k_{3x}^{\perp q} \text{Re } Y_n^1 R_n^1 (z P_J'') \right], \\ V_2^{(J)} &= \sum_{n=0}^J C_2^{(J)} k_{3z}^{\perp q} \left[\frac{2}{3} k_{3z}^{\perp q} Y_n^0 R_n^0 (z (P_J' + 2z P_J'')) \right. \\ &\quad \left. + k_{3x}^{\perp q} \text{Re } Y_n^1 R_n^1 (P_J' + 2z P_J'') \right], \end{aligned} \quad (35)$$

where

$$\begin{aligned} R_n^0(f) &= \int \frac{d\Omega}{4\pi} f(z) Y_n^0(z, \Theta), \\ R_n^1(f) &= 2 \int \frac{d\Omega}{4\pi} f(z) \cos \varphi \sin \Theta \text{Re } Y_n^1(z, \Theta). \end{aligned} \quad (36)$$

The P -vector amplitudes for the reggeized π_2 -exchanges read:

$$\begin{aligned} A_{-2}^{(J)}(s) &= \hat{P}_{-2}^{(J)} (I - i\hat{\rho}^J(s) \hat{K}^J)^{-1}, \\ A_0^{(J)}(s) &= \hat{P}_0^{(J)} (I - i\hat{\rho}^J(s) \hat{K}^J)^{-1}. \end{aligned} \quad (37)$$

The P -vector components are parametrized in the form:

$$\begin{aligned}
(P_{-2}^{(J)})_n &= \sum_{\alpha} \frac{1}{B_{J-2}(-p_{\perp}^2, r_{\alpha})} \left(\frac{G_{-2}^{(J)\alpha} g_n^{\alpha(J)}}{M_{\alpha}^2 - s} \right) \frac{1}{B_J(-k_{n\perp}^2, r_{\alpha})} \\
&+ \frac{F_{(-2)n}^{(J)}}{B_{J-2}(-p_{\perp}^2, r_0) B_J(-k_{n\perp}^2, r_0)}, \\
(P_{(0)}^{(J)})_n &= \sum_{\alpha} \frac{1}{B_J(-p_{\perp}^2, r_{\alpha})} \left(\frac{G_0^{(J)\alpha} g_n^{\alpha(J)}}{M_{\alpha}^2 - s} \right) \frac{1}{B_J(-k_{n\perp}^2, r_{\alpha})} \\
&+ \frac{F_{(0)n}^{(J)}}{B_J(-p_{\perp}^2, r_0) B_J(-k_{n\perp}^2, r_0)}
\end{aligned} \tag{38}$$

The total amplitude of the π_2 exchange can be rewritten as an expansion over spherical functions:

$$A_{\pi p \rightarrow \pi \pi n}^{(\pi_2)} = \sum_{n=0}^N \left(Y_n^0 A_{tot}^{0(n)}(s) + Y_n^1 A_{tot}^{1(n)}(s) \right) R_{\pi_2}(s_{\pi N}, q^2) \left(\varphi_n^+(\vec{\sigma} \vec{p}_{\perp}) \varphi_p \right) g_{pn}^{(\pi_2)}, \tag{39}$$

where

$$\begin{aligned}
A_{tot}^{0(n)}(s) &= \frac{1}{s_{\pi N}^2} (k_{3z}^{\perp q})^2 \sum_J \left[R_n^0(P_2 P_J'') C_3^{(J)} A_{-2}^{(J)}(s) \right. \\
&\quad \left. - \frac{2}{3} R_n^0(z(P_J' + 2z P_J'')) C_2^{(J)} A_0^{(J)}(s) \right], \\
A_{tot}^{1(n)}(s) &= \frac{1}{s_{\pi N}^2} k_{3z}^{\perp q} k_{3x} \sum_J \left[3 R_n^1(P_2 P_J'') C_3^{(J)} A_{-2}^{(J)}(s) \right. \\
&\quad \left. - R_n^1(z(P_J' + 2z P_J'')) C_2^{(J)} A_0^{(J)}(s) \right].
\end{aligned} \tag{40}$$

Then the final expression is:

$$\begin{aligned}
N(M, t) \langle Y_J^0 \rangle &= \frac{\rho(s) \sqrt{s}}{\pi |\vec{p}_2|^2 s_{\pi N}} R_{\pi_2}(s_{\pi N}, q^2) R_{\pi_2}^*(s_{\pi N}, q^2) (-t) (g_{pn}^{(\pi_2)})^2 \\
&\times \sum_{n,m} \left[d_{n,m,J}^{000} A_{tot}^{0(n)}(s) A_{tot}^{0(m)*}(s) + d_{n,m,J}^{110} A_{tot}^{1(n)}(s) A_{tot}^{1(m)*}(s) \right], \\
N(M, t) \langle Y_J^1 \rangle &= \frac{\rho(s) \sqrt{s}}{\pi |\vec{p}_2|^2 s_{\pi N}} R_{\pi_2}(s_{\pi N}, q^2) R_{\pi_2}^*(s_{\pi N}, q^2) (-t) (g_{pn}^{(\pi_2)})^2 \\
&\times \sum_{n,m} \left[d_{n,m,J}^{101} A_{tot}^{1(n)}(s) A_{tot}^{0(m)*}(s) + d_{n,m,J}^{011} A_{tot}^{0(n)}(s) A_{tot}^{1(m)*}(s) \right].
\end{aligned} \tag{41}$$

3 Status of trajectories on (J, M^2) plane

The π , η , a_1 , a_2 , a_3 , ρ and P' (or f_2) trajectories on (J, M^2) planes are shown in Fig. 1.

Leading π and η trajectories are unambiguously determined together with their daughter trajectories, while for a_2 , a_1 , ρ and P' only the leading trajectories can be given in a definite way.

In the construction of (J, M^2) -trajectories it is essential that the leading meson trajectories (π , ρ , a_1 , a_2 and P') are well known from the analysis of the diffraction scattering of hadrons at $p_{lab} \sim 5 - 50$ GeV/c (for example, see [3] and references therein).

The pion and η trajectories are linear with a good accuracy (see Fig. 1). Other leading trajectories (ρ , a_1 , a_2 , P') can also be considered as linear:

$$\alpha_X(M^2) \simeq \alpha_X(0) + \alpha'_X(0)M^2. \quad (42)$$

The parameters of the linear trajectories, determined by the masses of the $q\bar{q}$ states, are

$$\begin{aligned} \alpha_\pi(0) &\simeq -0.015, & \alpha'_\pi(0) &\simeq 0.83 \text{ GeV}^{-2}; \\ \alpha_\rho(0) &\simeq 0.50, & \alpha'_\rho(0) &\simeq 0.87 \text{ GeV}^{-2}; \\ \alpha_\eta(0) &\simeq -0.25, & \alpha'_\eta(0) &\simeq 0.80 \text{ GeV}^{-2}; \\ \alpha_{a_1}(0) &\simeq -0.10, & \alpha'_{a_1}(0) &\simeq 0.72 \text{ GeV}^{-2}; \\ \alpha_{a_2}(0) &\simeq 0.45, & \alpha'_{a_2}(0) &\simeq 0.93 \text{ GeV}^{-2}; \\ \alpha_{P'}(0) &\simeq 0.50, & \alpha'_{P'}(0) &\simeq 0.93 \text{ GeV}^{-2}. \end{aligned} \quad (43)$$

The slopes $\alpha'_X(0)$ of the trajectories are approximately equal. The inverse slope, $1/\alpha'_X(0) \simeq 1.25 \pm 0.15 \text{ GeV}^2$, roughly equals the parameter μ^2 for trajectories on the (n, M^2) planes:

$$\frac{1}{\alpha'_X(0)} \simeq \mu^2. \quad (44)$$

In the subsequent chapters, considering the scattering processes, we use for the Regge trajectories the momentum transfer squared $M^2 \rightarrow t$.

3.1 Kaon trajectories on (J, M^2) plane

As was said above, experimental data in the kaon sector are scarce, so in Fig. 2 we show only the leading K -meson trajectory (the states with $J^P = 0^-, 2^-$), the K^* trajectory ($J^P = 1^-, 3^-, 5^-$) and the leading and daughter trajectories for $J^P = 0^+, 2^+, 4^+$. The parameters of the leading kaon trajectories are as follows:

$$\begin{aligned} \alpha_K(0) &\simeq -0.25, & \alpha'_K(0) &\simeq 0.90 \text{ GeV}^{-2}; \\ \alpha_{K^*}(0) &\simeq 0.30, & \alpha'_{K^*}(0) &\simeq 0.85 \text{ GeV}^{-2}; \\ \alpha_{K_{2^+}}(0) &\simeq -0.2, & \alpha'_{K_{2^+}}(0) &\simeq 1.0 \text{ GeV}^{-2}. \end{aligned} \quad (45)$$




Figure 1: Trajectories in the (J, M^2) plane: a) leading and daughter π -trajectories, b) leading and daughter η -trajectories, c) a_2 -trajectories, d) leading and daughter a_1 -trajectories, e) ρ -trajectories, f) P' -trajectories.




Figure 2: Kaon trajectories on the (J^P, M^2) plane.

The trajectories with $J^P = 1^+, 3^+, 5^+$ cannot be defined unambiguously.

4 Dispersion integral equation for a three-body system and the K -matrix approach

The LEAR (CERN) experiment accumulated high statistics data on three-meson production from the $\bar{p}p$ annihilation at rest, mainly from $(J^{PC} = 0^{-+})$ -level. The data of the Crystal Barrel Collaboration (LEAR) were successfully analysed in the terms of the K -matrix approach (see, for example, [4, 11, 12, 15]) with the aim to search for new meson resonances in the region 1000–1600 MeV. But the K -matrix approach takes into account three-body unitarity condition on the phenomenological level only. The three-body unitarity can be taken into account considering all three-particle rescatterings, using, for example, the dispersion relation N/D -method.

In this Appendix we demonstrate what type of assumptions is needed for transformation of the three-body dispersion relation amplitudes into that of the K -matrix approach.

Here the dispersion relation N/D -method is presented for a three-body system: the method allows one to take into account final-state two-meson interactions. We consider in detail an illustrative example: the decay of the 0^{-+} -state into three different pseudoscalar mesons.




Figure 3: Different types of transitions ($J_{in}^P = 0^-$)-state $\longrightarrow P_1 P_2 P_3$: a) prompt decay, b) decay with subsequent final state interactions.

The first steps in accounting for all two-body final state interactions were made in [16] in a non-relativistic approach for three-nucleon systems. In [17] the two-body interactions were considered in the potential approach (the Faddeev equation).

The relativistic dispersion relation technique was used for the investigation of the final state interaction effects in [18].

A relativistic dispersion relation equation for the amplitude $\eta \rightarrow \pi\pi\pi$ was written in [19]. Later on the method was generalised [20] for the coupled processes $p\bar{p}(\text{at rest}) \rightarrow \pi\pi\pi, \eta\eta\pi, K\bar{K}\pi$: this way a system of coupled equations for decay amplitudes was written. Following [19, 20], we explain here the main points in considering the dispersion relations for a three-particle system. The account of the three-particle final state interactions imposes correct unitarity and analyticity constraints on the amplitude.

4.0.1 Two-particle interactions in the 0^- -state $\longrightarrow P_1 P_2 P_3$ decay

As previously, we consider the decay of a pseudoscalar particle ($J_{in}^P = 0^-$) with the mass M and momentum P into three pseudoscalar particles with masses m_1, m_2, m_3 and momenta k_1, k_2, k_3 . There are different contributions to this decay process: those without final state particle interactions (prompt decay, Fig. 3a) and decays with subsequent final state interactions (an example is shown in Fig. 3b).

For the decay amplitude we consider here an equation which takes into account two-particle final state interactions, such as that shown in Fig. 3b. First, we consider in detail the S -wave interactions. This case clarifies the main points of the dispersion relation approach for the three-particle interaction amplitude. Then we discuss a scheme for generalising the equations for the case of higher waves.

(i) S -wave interaction.

Let us begin with the S -wave two-particle interactions. The decay amplitude is given by

$$A_{P_1 P_2 P_3}^{(J_{in}=0)}(s_{12}, s_{13}, s_{23}) = \lambda(s_{12}, s_{13}, s_{23}) + A_{12}^{(0)}(s_{12}) + A_{13}^{(0)}(s_{13}) + A_{23}^{(0)}(s_{23}). \quad (46)$$

Different terms in (46) are illustrated by Fig. 4: we have a prompt production amplitude, Fig. 4b, and terms $A_{ij}^{(0)}(s_{ij})$ with particles $P_1 P_2$ (Fig. 4c), $P_1 P_3$ (Fig. 4d) and $P_2 P_3$ (Fig. 4e) participating in final state interactions.




Figure 4: Different terms in the amplitude $A_{P_1 P_2 P_3}^{(J_{in}=0)}(s_{12}, s_{13}, s_{23})$

To take into account rescattering of the type shown in Fig. 3b, we can write equations for different terms $A_{ij}^{(0)}(s_{ij})$.

The two-particle unitarity condition is explored to derive the integral equation for the amplitude $A_{ij}^{(0)}(s_{ij})$. The idea of the approach suggested in [21] is that one should consider the case of a small external mass $M < m_1 + m_2 + m_3$. A standard spectral integral equation (or a dispersion relation equation) is written in this case for the transitions $h_{in} P_\ell \rightarrow P_i P_j$. Then the analytical continuation is performed over the mass M back to the decay region: this gives a system of equations for decay amplitudes $A_{ij}^{(0)}(s_{ij})$.

So, let us consider the channel of particles 1 and 2, the transition $h_{in} P_3 \rightarrow P_1 P_2$. We write the two-particle unitarity condition for the scattering in this channel with the assumption $(M + m_3) \sim (m_1 + m_2)$.

The discontinuity of the amplitude in the s_{12} -channel equals

$$\begin{aligned} disc_{12} A_{P_1 P_2 P_3}^{J_{in}=0}(s_{12}, s_{13}, s_{23}) &= disc_{12} A_{12}^{(0)}(s_{12}) = \\ &= \int d\Phi_{12}(p_{12}; k_1, k_2) \left(\lambda(s_{12}, s_{13}, s_{23}) + A_{12}^{(0)}(s_{12}) + A_{13}^{(0)}(s_{13}) + A_{23}^{(0)}(s_{23}) \right) \left(A_{12 \rightarrow 12}^{(0)}(s_{12}) \right)^* . \end{aligned} \quad (47)$$

Here $d\Phi_{12}(p_{12}; k_1, k_2) = (1/2)(2\pi)^{-2} \delta^4(p_{12} - k_1 - k_2) d^4 k_1 d^4 k_2 \delta(m_1^2 - k_1^2) \delta(m_2^2 - k_2^2)$ is the standard phase volume of particles 1 and 2. In (47), we should take into account that only $A_{12}^{(0)}(s_{12})$ has a non-zero discontinuity in the channel 12.

S-wave two-particle scattering amplitude

But first, let us consider the S -wave two-particle scattering amplitude $A_{P_1 P_2 \rightarrow P_1 P_2}^{(0)}$. It can be written in the dispersion N/D approach with separable interaction (see Chapter 3) as a series

$$\begin{aligned} A_{P_1 P_2 \rightarrow P_1 P_2}^{(0)}(s) &= G_0^L(s_{12}) G_0^R(s_{12}) + G_0^L(s_{12}) B_{12}^{(0)}(s_{12}) G_0^R(s_{12}) \\ &+ G_0^L(s_{12}) B_{12}^{(0)2}(s_{12}) G_0^R(s_{12}) + \dots = \frac{G_0^L(s_{12}) G_0^R(s_{12})}{1 - B_{12}^{(0)}(s_{12})}, \end{aligned} \quad (48)$$

where $G_0^L(s_{12})$ and $G_0^R(s_{12})$ are left and right vertex functions. The loop diagram $B_{12}^{(0)}(s_{12})$ in

the dispersion relation representation reads:

$$B_{12}^{(0)}(s_{12}) = \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \frac{G_0^L(s'_{12})\rho_{12}^{(0)}(s'_{12})G_0^R(s'_{12})}{s'_{12} - s_{12} - i0}, \quad (49)$$

where $\rho_{12}^{(0)}(s_{12}) = \sqrt{[s_{12} - (m_1 + m_2)^2][s_{12} - (m_1 - m_2)^2]}/(16\pi s_{12})$ is the two-particle S -wave phase space integrated over the angular variables. The vertex functions contain left-hand singularities related to the t -channel exchange diagrams, while the loop diagram $B_{12}^{(0)}(s_{12})$ has a singularity due to the elastic scattering (the right-hand side singularity). The consideration of the scattering amplitude $A_{P_1 P_2 \rightarrow P_1 P_2}^{(0)}(s_{12})$ does not specify it whether both vertices, $G_0^L(s_{12})$ and $G_0^R(s_{12})$, have left-hand singularities or only one of them (see discussion in Chapter 3). Considering the three-body decay, it is convenient to make use of this freedom. On the first sheet of the decay amplitude, we take into account the threshold singularities at $s_{ij} = (m_i + m_j)^2$, which are associated with the elastic scattering in the subchannel of particles i and j but not those on the left-hand side. This means that the vertex $G_0^R(s_{12})$ should be chosen here as an analytical function. For the sake of simplicity let us put $G_0^R(s_{12}) = 1$ and present the amplitude $P_1 P_2 \rightarrow P_1 P_2$ as

$$A_{P_1 P_2 \rightarrow P_1 P_2}^{(0)}(s_{12}) = G_0^L(s_{12}) \frac{1}{1 - B_{12}^{(0)}(s_{12})} \quad \text{at} \quad G_0^R(s_{12}) = 1. \quad (50)$$

Equation for the decay amplitude $h_{in} \rightarrow P_1 P_2 P_3$

Exploring (47), let us now return to the equation for the decay amplitude $h_{in} \rightarrow P_1 P_2 P_3$.

As was noted (see also [21]), the full set of rescattering of particles 1 and 2 gives us the factor $(1 - B_0(s_{12}))^{-1}$, so we have from (47):

$$A_{12}^{(0)}(s_{12}) = B_{in}^{(0)}(s_{12}) \frac{1}{1 - B_0(s_{12})}. \quad (51)$$

The first loop diagram $B_{in}^{(0)}(s_{12})$ is determined as

$$B_{in}^{(0)}(s_{12}) = \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \frac{disc_{12} B_{in}^{(0)}(s'_{12})}{s'_{12} - s_{12} - i0}, \quad (52)$$

where

$$\begin{aligned} disc_{12} B_{in}^{(0)}(s_{12}) &= \int d\Phi_{12}(p_{12}; k_1, k_2) \left(\lambda(s_{12}, s_{13}, s_{23}) + A_{13}^{(0)}(s_{13}) + A_{23}^{(0)}(s_{23}) \right) G_0^L(s_{12}) \\ &\equiv disc_{12} B_{\lambda-12}^{(0)}(s_{12}) + disc_{12} B_{13-12}^{(0)}(s_{12}) + disc_{12} B_{23-12}^{(0)}(s_{12}). \end{aligned} \quad (53)$$

Here we present $disc_{12} B_{in}^{(0)}(s_{12})$ as a sum of three terms because each of them needs a special treatment when $M^2 + i\varepsilon$ is increasing.

It is convenient to perform the phase-space integration in equation (53) in the centre-of-mass system of particles 1 and 2 where $\mathbf{k}_1 + \mathbf{k}_2 = 0$. In this frame

$$\begin{aligned} s_{13} &= m_1^2 + m_3^2 + 2k_{10}k_{30} - 2z |\mathbf{k}_1| |\mathbf{k}_3|, \\ s_{23} &= m_2^2 + m_3^2 + 2k_{20}k_{30} + 2z |\mathbf{k}_2| |\mathbf{k}_3|, \end{aligned} \quad (54)$$

where $z = \cos \theta_{13}$ and $k_{10} = \frac{s_{12} + m_1^2 - m_2^2}{2\sqrt{s_{12}}}$, $k_{20} = \frac{s_{12} + m_2^2 - m_1^2}{2\sqrt{s_{12}}}$, $-k_{30} = \frac{s_{12} + m_3^2 - M^2}{2\sqrt{s_{12}}}$. The minus sign in front of k_{30} reflects the fact that P_3 is an outgoing, not an incoming particle. As usually, $|\mathbf{k}_j| = \sqrt{k_{j0}^2 - m_j^2}$ for $j = 1, 2, 3$. In the calculation of $disc_{12} B_0^{(in)}(s_{12})$ all integrations are carried out easily except for the contour integral over dz . It can be rewritten in (53) as an integral over ds_{13} or ds_{23} :

$$\int_{-1}^{+1} \frac{dz}{2} \rightarrow \int_{s_{13}(-)}^{s_{13}(+)} \frac{ds_{13}}{4 |\mathbf{k}_1| |\mathbf{k}_3|}, \quad \text{or} \quad \int_{-1}^{+1} \frac{dz}{2} \rightarrow \int_{s_{23}(-)}^{s_{23}(+)} \frac{ds_{23}}{4 |\mathbf{k}_2| |\mathbf{k}_3|}, \quad (55)$$

where

$$\begin{aligned} s_{13}(\pm) &= m_1^2 + m_3^2 + 2k_{10}k_{30} \pm 2 |\mathbf{k}_1| |\mathbf{k}_3|, \\ s_{23}(\pm) &= m_2^2 + m_3^2 + 2k_{20}k_{30} \pm 2 |\mathbf{k}_2| |\mathbf{k}_3|. \end{aligned} \quad (56)$$

The relative location of the integration contours (55) and amplitude singularities is the determining point for writing the equation.

Below we use the notation

$$\int_{s_{i3}(-)}^{s_{i3}(+)} ds_{i3} = \int_{C_{i3}(s_{12})} ds_{i3}. \quad (57)$$

One can see from (56) that the integration contours $C_{13}(s_{12})$ and $C_{23}(s_{12})$ depend on M^2 and s_{12} , so we should monitor them when $M^2 + i\varepsilon$ increases.

Let us underline again that the idea to consider the decay processes in the dispersion relation approach is the following : we write the equation in the region of the standard scattering *two particles* $\rightarrow$ *two particles* (when $m_1 \sim m_2 \sim m_3 \sim M$) with the subsequent analytical continuation (with $M^2 + i\varepsilon$ at $\varepsilon > 0$) into the decay region, $M > m_1 + m_2 + m_3$, and then $\varepsilon \rightarrow +0$. In this continuation we need to specify what type of singularities (and corresponding type of processes) we take into account and what type of singularities we neglect. Definitely, we take into account right-hand side and left-hand side singularities of the scattering processes $P_i P_j \rightarrow P_i P_j$ (our main aim is to restore the rescattering processes correctly). But singularities of the prompt production amplitude are beyond the field of our interest. In other words, we suppose $\lambda(s_{12}, s_{13}, s_{23})$ to be an analytical function in the region under consideration.

Assuming $\lambda(s_{12}, s_{13}, s_{23})$ to be an analytical function in the region under consideration, we can easily perform analytical continuation of the integral over dz , Eq. (55), with $M^2 + i\varepsilon$.

Problems may appear in the integrations of $A_{13}^{(0)}(s_{13})$ and $A_{23}^{(0)}(s_{23})$ owing to the threshold singularities in the amplitudes (at $s_{13} = (m_1 + m_3)^2$ and $s_{23} = (m_2 + m_3)^2$, respectively). However, the analytical continuation over $M^2 + i\varepsilon$ resolves them: one can see in [2] (Chapter 4) the location of the integration contour in the complex- s_{23} plane with respect to the threshold singularity at $s_{23} = (m_2 + m_3)^2$ when $M > m_1 + m_2 + m_3$.

Let us now write the equation for the three-particle production amplitude in more detail. We denote the S -wave projection of $\lambda(s_{12}, s_{13}, s_{23})$ as

$$\langle \lambda(s_{12}, s_{13}, s_{23}) \rangle_{12}^{(0)} = \int_{-1}^{+1} \frac{dz}{2} \lambda(s_{12}, s_{13}, s_{23}), \quad (58)$$

and the contour integrals over the amplitudes $A_{13}^{(0)}(s_{13})$ and $A_{23}^{(0)}(s_{23})$ as

$$\langle A_{i3}^{(0)}(s_{i3}) \rangle_{12}^{(0)} = \int_{-1}^{+1} \frac{dz}{2} A_{i3}^{(0)}(s_{i3}) \equiv \int_{C_{i3}(s_{12})} \frac{ds_{i3}}{4|\mathbf{k}_i||\mathbf{k}_3|} A_{i3}^{(0)}(s_{i3}), \quad i = 1, 2. \quad (59)$$

Remind once more that the definition of the contours $C_i(s_{12})$ is given in (57) while the relative position of the contour $C_2(s_{12})$ and the threshold singularity in the s_{23} -channel is shown in Fig. 4.26. So, we rewrite (53) in the form

$$\begin{aligned} disc_{12} B_{in}^{(0)}(s_{12}) &= \left(\langle \lambda(s_{12}, s_{13}, s_{23}) \rangle_{12}^{(0)} + \langle A_{13}^{(0)}(s_{13}) \rangle_{12}^{(0)} + \langle A_{23}^{(0)}(s_{23}) \rangle_{12}^{(0)} \right) \\ &\times \rho_{12}^{(0)}(s_{12}) G_0^L(s_{12}), \end{aligned} \quad (60)$$

Equation (60) allows us to write the dispersion integral for the loop amplitude $B_{in}^{(0)}(s_{12})$. As a result, we have:

$$A_{12}^{(0)}(s_{12}) = \left(B_{\lambda-12}^{(0)}(s_{12}) + B_{13-12}^{(0)}(s_{12}) + B_{23-12}^{(0)}(s_{12}) \right) \frac{1}{1 - B_{12}^{(0)}(s_{12})} \quad (61)$$

where

$$\begin{aligned} B_{\lambda-12}^{(0)}(s_{12}) &= \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(0)} \frac{\rho_{12}^{(0)}(s'_{12})}{s'_{12} - s_{12} - i0} G_0^L(s'_{12}), \\ B_{i3-12}^{(0)}(s_{12}) &= \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \langle A_{i3}^{(0)}(s'_{i3}) \rangle_{12}^{(0)} \frac{\rho_{12}^{(0)}(s'_{12})}{s'_{12} - s_{12} - i0} G_0^L(s'_{12}). \end{aligned} \quad (62)$$

Let us emphasise that in the integrand (62) the energy squared is s'_{12} and hence, calculating $\langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(0)}$ and $\langle A_{i3}^{(0)}(s'_{i3}) \rangle_{12}^{(0)}$, we should use Eqs. (54) – (57) with the replacement $s_{12} \rightarrow s'_{12}$.

The equation (61) is illustrated by Fig. 5.




Figure 5: Diagrammatic presentation of Eq. (61).

In the same way we can write equations for $A_{13}^{(0)}(s_{13})$ and $A_{23}^{(0)}(s_{23})$. We have a system of three non-homogeneous equations which determine the amplitudes $A_{ij}^{(0)}(s_{ij})$ when $\lambda(s_{12}, s_{13}, s_{23})$ is considered as an input function.

Note that the integration contour $C_i(s_{12})$ in (59), see also [21], does not coincide with that of [22] where the corresponding problem was treated starting from the consideration of the three-body channel.

(ii) Final state rescattering $P_i P_j \rightarrow P_i P_j$ in the $L > 0$ state.

Equations for amplitudes which describe the final state interactions in the transition $(J_{in}^P = 0^-)$ -state $\rightarrow P_1 P_2 P_3$ when rescattering $P_i P_j \rightarrow P_i P_j$ occur in a state with $L > 0$ can be written in a way analogous to that presented above for $L = 0$. So, we suppose that $P_i P_j \rightarrow P_i P_j$ rescattering take place in a state with definite orbital momentum L and $L \neq 0$.

The amplitude for the decay $(J_{in}^P = 0^-)$ -state $\rightarrow P_1 P_2 P_3$ (below $L = J$) reads:

$$A_{P_1 P_2 P_3}^{(J_{in}=0)}(s_{12}, s_{13}, s_{23}) = \lambda(s_{12}, s_{13}, s_{23}) + A_{12}^{(J)}(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp P}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}) \\ + A_{13}^{(J)}(s_{13}) X_{\mu_1 \dots \mu_J}^{(J)}(k_2^{\perp P}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{13}^{\perp p_{13}}) + A_{23}^{(J)}(s_{23}) X_{\mu_1 \dots \mu_J}^{(J)}(k_1^{\perp P}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{23}^{\perp p_{23}}). \quad (63)$$

Convolutions of the momentum operators, such as $X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp P}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}})$, being functions of s_{ij} do not contain threshold singularities. So we can rewrite (63) in a more compact form

$$A_{P_1 P_2 P_3}^{(J_{in}=0)}(s_{12}, s_{13}, s_{23}) = \lambda(s_{12}, s_{13}, s_{23}) + A_{12}^{(J-J)}(s_{12}, s_{13}, s_{23}) \\ + A_{13}^{(J-J)}(s_{12}, s_{13}, s_{23}) + A_{23}^{(J-J)}(s_{12}, s_{13}, s_{23}), \quad (64)$$

where $A_{12}^{(J-J)}(s_{12}, s_{13}, s_{23}) = A_{12}^{(J)}(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp P}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}})$, and so on. As previously, $\lambda(s_{12}, s_{13}, s_{23})$ is an analytical function of s_{ij} while the terms $A_{ij}^{(J-J)}(s_{12}, s_{13}, s_{23})$ have threshold singularities of the type $\sqrt{s_{ij} - (m_i + m_j)^2}$ due to final state rescattering $P_i P_j \rightarrow P_i P_j$.

Two-particle scattering amplitude at $L = J \neq 0$

First, we should introduce a two-particle ($L = J$)-wave scattering amplitude. To be definite, we consider $P_1 P_2 \rightarrow P_1 P_2$. We write the block of a one-fold scattering as

$$\begin{aligned} A_{(P_1 P_2 \rightarrow P_1 P_2)_{\text{one-fold}}}^{(J)}(s_{12}) &= X_{\nu_1 \dots \nu_J}^{(J)}(k_{12}^{\perp p_{12}}) O_{\mu_1 \dots \mu_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) G_J^L(s_{12}) G_J^R(s_{12}) \\ &\times X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}). \end{aligned} \quad (65)$$

The two-fold scattering amplitude reads:

$$\begin{aligned} A_{(P_1 P_2 \rightarrow P_1 P_2)_{\text{two-fold}}}^{(J)}(s_{12}) &= X_{\nu_1 \dots \nu_J}^{(J)}(k_{12}^{\perp p_{12}}) O_{\nu'_1 \dots \nu'_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) G_J^L(s_{12}) \\ &\times \left[\int_{(m_1+m_2)^2}^{\infty} \frac{ds''_{12}}{\pi(s''_{12} - s_{12} - i0)} G_J^R(s''_{12}) \right. \\ &\times \left. \int X_{\nu'_1 \dots \nu'_J}^{(J)}(k_{12}^{\perp p''_{12}}) d\Phi_{12}(p''_{12}; k''_1, k''_2) X_{\nu''_1 \dots \nu''_J}^{(J)}(k_{12}^{\perp p''_{12}}) G_J^L(s''_{12}) \right] \\ &\times G_J^R(s_{12}) O_{\mu_1 \dots \mu_J}^{\nu''_1 \dots \nu''_J}(\perp p_{12}) G_J^L(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}). \end{aligned} \quad (66)$$

In the integrand we can replace $k_{12}^{\perp p''_{12}} \rightarrow k_{12}^{\perp p_{12}}$, because in the c.m. frame of particles $P_1 P_2$ one has $p''_{12} = (\sqrt{s''_{12}}, 0, 0, 0)$ and $p_{12} = (\sqrt{s_{12}}, 0, 0, 0)$. The integration over the phase space gives

$$\begin{aligned} \int X_{\nu'_1 \dots \nu'_J}^{(J)}(k_{12}^{\perp p_{12}}) d\Phi_{12}(p''_{12}; k''_1, k''_2) X_{\nu''_1 \dots \nu''_J}^{(J)}(k_{12}^{\perp p_{12}}) &= O_{\nu'_1 \dots \nu'_J}^{\nu''_1 \dots \nu''_J}(\perp p_{12}) \\ &\times \rho_{12}^{(J)}(s''_{12}). \end{aligned} \quad (67)$$

Using

$$O_{\nu'_1 \dots \nu'_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) O_{\nu''_1 \dots \nu''_J}^{\nu'_1 \dots \nu'_J}(\perp p_{12}) O_{\mu_1 \dots \mu_J}^{\nu''_1 \dots \nu''_J}(\perp p_{12}) = O_{\mu_1 \dots \mu_J}^{\nu_1 \dots \nu_J}(\perp p_{12}), \quad (68)$$

we write the two-fold amplitude as follows:

$$\begin{aligned} A_{(P_1 P_2 \rightarrow P_1 P_2)_{\text{two-fold}}}^{(J)}(s_{12}) &= \\ &X_{\nu_1 \dots \nu_J}^{(J)}(k_{12}^{\perp p_{12}}) O_{\mu_1 \dots \mu_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) G_J^L(s_{12}) B_{12}^{(J)}(s_{12}) G_J^R(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}), \end{aligned} \quad (69)$$

where

$$B_{12}^{(J)}(s_{12}) = \int_{(m_1+m_2)^2}^{\infty} \frac{ds''_{12}}{\pi(s''_{12} - s_{12} - i0)} G_J^R(s''_{12}) \rho_{12}^{(J)}(s''_{12}) G_J^L(s''_{12}) \quad (70)$$

is the loop diagram.

The full set of rescattering gives:

$$\begin{aligned} A_{P_1 P_2 \rightarrow P_1 P_2}^{(J)}(s_{12}) &= X_{\nu_1 \dots \nu_J}^{(J)}(k_{12}^{\perp p_{12}}) O_{\mu_1 \dots \mu_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) \\ &\times G_J^L(s_{12}) \frac{1}{1 - B_{12}^{(J)}(s_{12})} G_J^R(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}). \end{aligned} \quad (71)$$

As previously for $J = 0$, we put

$$G_J^R(s_{12}) = 1. \quad (72)$$

Finally we write:

$$A_{P_1 P_2 \rightarrow P_1 P_2}^{(J)}(s_{12}) = X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}) G_J^L(s_{12}) \frac{1}{1 - B_{12}^{(J)}(s_{12})} X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}). \quad (73)$$

Equation for the decay amplitude $h_{J_{in}=0} \rightarrow P_1 P_2 P_3$

Let us return now to the equation for the three-particle production amplitude $A_{P_1 P_2 P_3}^{(J_{in}=0)}(s_{12}, s_{13}, s_{23})$ given by (64). We write equations for separated terms $A_{ij}^{(J-J)}(s_{12}, s_{13}, s_{23})$. To use Eqs. (65)–(73) directly, we consider the term with the final state interaction in the channel 12, namely, $A_{12}^{(J-J)}(s_{12}, s_{13}, s_{23})$.

The amplitude $A_{12}^{(J-J)}(s_{12}, s_{13}, s_{23})$ is determined by three terms shown in Fig. 5b,c,d.

The term initiated by the prompt production block $\lambda(s_{12}, s_{13}, s_{23})$ is a set of loop diagrams of the type of that in Fig. 5b. Therefore this term reads:

$$X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp p_{12}}) B_{\lambda-12}^{(J)}(s_{12}) \frac{1}{1 - B_{12}^{(J)}(s_{12})} X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}), \quad (74)$$

with

$$B_{\lambda-12}^{(J)}(s_{12}) = \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J)} \frac{\rho_{12}^{(J)}(s'_{12})}{s'_{12} - s_{12} - i0} G_0^L(s'_{12}). \quad (75)$$

Let us explain Eqs. (74), (75) in more detail. Similarly to (65), we write for the first loop diagram in (74) the following representation:

$$\begin{aligned} & X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp p_{12}}) B_{\lambda-12}^{(J)}(s_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}) = X_{\nu_1 \dots \nu_J}^{(J)}(k_3^{\perp p_{12}}) O_{\nu'_1 \dots \nu'_J}^{\nu_1 \dots \nu_J}(\perp p_{12}) \\ & \times \left[\int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \frac{\langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J)}}{(s'_{12} - s_{12} - i0)} \int X_{\nu'_1 \dots \nu'_J}^{(J)}(k_{12}^{\perp p_{12}}) d\Phi_{12}(p'_{12}; k'_1, k'_2) X_{\nu'_1 \dots \nu'_J}^{(J)}(k_{12}^{\perp p_{12}}) G_J^L(s'_{12}) \right] \\ & \times O_{\mu_1 \dots \mu_J}^{\nu'_1 \dots \nu'_J}(\perp p_{12}) X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}}). \end{aligned} \quad (76)$$

Recall that $\langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J)}$ depends on s'_{12} only. Indeed, the projection $\langle \lambda(s_{12}, s_{13}, s_{23}) \rangle_{12}^{(J)}$ is determined by the following expansion of the non-singular term:

$$\lambda(s_{12}, s_{13}, s_{23}) = \sum_{J'} X_{\nu_1 \dots \nu_{J'}}^{(J')} (k_3^{\perp p_{12}}) \langle \lambda(s_{12}, s_{13}, s_{23}) \rangle_{12}^{(J')} X_{\mu_1 \dots \mu_{J'}}^{(J')} (k_{12}^{\perp p_{12}}), \quad (77)$$

so we have

$$\begin{aligned}
\langle \lambda(s_{12}, s_{13}, s_{23}) \rangle_{12}^{(J)} &= \int d\Phi_{h_{in}3}(p_{12}; P, -k_3) X_{\nu_1 \dots \nu_J}^{(J)}(k_3^{\perp p_{12}}) \lambda(s_{12}, s_{13}, s_{23}) \\
&\times X_{\mu_1 \dots \mu_{J'}}^{(J)}(k_{12}^{\perp p_{12}}) d\Phi_{12}(p_{12}; k_1, k_2) \int d\Phi_{h_{in}3}(p_{12}; P, -k_3) \left(X_{\nu'_1 \dots \nu'_{J'}}^{(J)}(k_3^{\perp p_{12}}) \right)^2 \\
&\times \int d\Phi_{12}(p_{12}; k_1, k_2) \left(X_{\mu'_1 \dots \mu'_{J'}}^{(J)}(k_{12}^{\perp p_{12}}) \right)^2.
\end{aligned} \tag{78}$$

In the integrand (76) the energy squared is s'_{12} . Hence, we should use $\langle \lambda(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(0)}$ in the calculation.

Likewise, we calculate the amplitudes of processes of Fig. 5c,d. As a result, we have the equation:

$$\begin{aligned}
A_{12}^{(J-J)}(s_{12}, s_{13}, s_{23}) &= X_{\mu_1 \dots \mu_J}^{(J)}(k_3^{\perp p_{12}}) \\
&\times \left(B_{\lambda-12}^{(J)}(s_{12}) + B_{13-12}^{(J)}(s_{12}) + B_{23-12}^{(J)}(s_{12}) \right) \frac{X_{\mu_1 \dots \mu_J}^{(J)}(k_{12}^{\perp p_{12}})}{1 - B_{12}^{(J)}(s_{12})}.
\end{aligned} \tag{79}$$

Here

$$B_{i3-12}^{(J)}(s_{12}) = \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \langle A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J)} \frac{\rho_{12}^{(J)}(s'_{12})}{s'_{12} - s_{12} - i0} G_0^L(s'_{12}). \tag{80}$$

Let us stress once more that in (80) the terms $\langle A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J')}$ and $\langle A_{23}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J')}$ depend on s'_{12} only. This is the result of the following expansions:

$$\begin{aligned}
A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) &= \sum_{J'} X_{\nu_1 \dots \nu_{J'}}^{(J')}(k_3^{\perp p_{12}}) \langle A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J')} X_{\mu_1 \dots \mu_{J'}}^{(J')}(k_{12}^{\perp p'_{12}}), \\
A_{23}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) &= \sum_{J'} X_{\nu_1 \dots \nu_{J'}}^{(J')}(k_3^{\perp p_{12}}) \langle A_{23}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{12}^{(J')} X_{\mu_1 \dots \mu_{J'}}^{(J')}(k_{12}^{\perp p'_{12}}).
\end{aligned} \tag{81}$$

The integration over the phase space in the calculations of $A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23})$ is performed in a way analogous to that for $J = 0$. The contour integrals read (see Fig. 6) :

$$\begin{aligned}
\langle A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \rangle_{i3}^{(J)} &= \int_{-1}^{+1} \frac{dz'}{2} A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}) \\
&\equiv \int_{C_{i3}(s'_{12})} \frac{ds_{i3}}{4|\mathbf{k}_i||\mathbf{k}_3|} A_{i3}^{(J-J)}(s'_{12}, s'_{13}, s'_{23}), \quad i = 1, 2.
\end{aligned} \tag{82}$$

The definition of the contours $C_i(s_{12})$ is given in (57).




Figure 6: The integration contour $C(s'_{12})$.

4.0.2 Dispersion relation equations for a three-body system with resonance interaction in the two-particle states of the outgoing hadrons

In this section we consider the case when the outgoing particles interact due to two-particle resonances. Such a situation occurs, for example, in the reaction $p\bar{p}(at\ rest, level\ ^1S_0) \rightarrow \pi\pi\pi$: in the 0^{++} -wave of the pion-pion amplitude, there is a set of comparatively narrow resonances while a non-resonance background can be described as a broad resonance. Another possibility to introduce the background contribution in this model is to add pole (resonance) terms beyond (for example, above) the region of application of the amplitude.

In order to avoid cumbersome formulas, we consider, as before, a reaction of the type $h(^1S_0) \rightarrow P_1P_2P_3$, with the S -wave interactions of the outgoing pseudoscalars $P_iP_j \rightarrow P_iP_j$.

(i) Two-particle resonance amplitude.

We start with the dispersion representation of the two-particle amplitude for this particular case. The first resonance term (the one-fold scattering block) of the amplitude $P_iP_j \rightarrow P_iP_j$ can be written in the form

$$\sum_n \frac{g_{ij}^{(n)2}(s_{ij})}{M_n^2 - s_{ij}}, \quad (83)$$

where M_n is a non-physical mass of the n -resonance, and vertex $g_{ij}^{(n)}(s_{ij})$ describes its decay into two particles P_iP_j . Experimental data tell us that vertices $g_{ij}^{(n)}(s_{ij})$ can be successfully approximated by the energy dependence: $g_{ij}^{(n)}(s_{ij}) \sim \exp(-s_{ij}/\mu^2)$ with the universal slope $\mu^2 \simeq 0.5\text{ GeV}^2$. Below we assume this universality:

$$g_{ij}^{(n)}(s_{ij}) = g_{ij}^{(n)} f(s_{ij}) \quad (84)$$

where $g_{ij}^{(n)}$ is a constant and $f(s_{ij})$ is a universal form factor of the type $\exp(-s_{ij}/\mu^2)$. If so, the two-fold scattering term of the amplitude contains the universal loop diagram $b(s_{ij})$:

$$A_{(P_iP_j \rightarrow P_iP_j)_{\text{two-fold}}}^{(0)}(s_{ij}) = f(s_{ij}) \sum_n \frac{g_{ij}^{(n)2}}{M_n^2 - s_{ij}} b(s_{ij}) \sum_{n'} \frac{g_{ij}^{(n')2}}{M_{n'}^2 - s_{ij}} f(s_{ij}),$$

$$b(s_{ij}) = \int_{(m_i+m_j)^2}^{\infty} \frac{ds' \rho_{ij}^{(0)}(s') f^2(s')}{\pi s' - s_{ij} - i0}. \quad (85)$$

Summing up the terms with different numbers of loops, one obtains the following expression for the amplitude:

$$A_{(P_i P_j \rightarrow P_i P_j)}^{(0)}(s_{ij}) = \frac{f^2(s_{ij}) \sum_n \frac{g_{ij}^{(n)2}}{M_n^2 - s_{ij}}}{1 - b(s_{ij}) \sum_{n'} \frac{g_{ij}^{(n')2}}{M_{n'}^2 - s_{ij}}}. \quad (86)$$

Since the loop diagram has the following real and imaginary parts:

$$\begin{aligned} b(s_{ij}) &= i \rho_{ij}^{(0)}(s_{ij}) f^2(s_{ij}) + P \int_{(m_i+m_j)^2}^{\infty} \frac{ds' \rho_{ij}^{(0)}(s') f^2(s')}{\pi s' - s_{ij}} \\ &= i \text{Im } b(s_{ij}) + \text{Re } b(s_{ij}), \end{aligned} \quad (87)$$

the scattering amplitude (86) can easily be rewritten in the K -matrix form for the case when an S -wave state contains several resonances.

(ii) Three particle production amplitude $h(^1S_0) \rightarrow P_1 P_2 P_3$.

The decay amplitude is given by an equation of the type of (46). In the term $\lambda(s_{12}, s_{13}, s_{23})$, however, we should take into account the prompt production of resonances; it is convenient to consider their widths as well. Correspondingly, we replace

$$\begin{aligned} \lambda(s_{12}, s_{13}, s_{23}) &\rightarrow \Lambda_{12}(s, s_{12}) + \Lambda_{13}(s, s_{13}) + \Lambda_{23}(s, s_{23}), \\ \Lambda_{ij}(s, s_{ij}) &= \sum_n \Lambda_{ij}^{(n)}(s, s_{ij}) \frac{g_{ij}^{(n)}}{M_n^2 - s_{ij}} f(s_{ij}) \frac{1}{1 - b(s_{ij}) \sum_{n'} \frac{g_{ij}^{(n')2}}{M_{n'}^2 - s_{ij}}}, \end{aligned} \quad (88)$$

and write for the full amplitude:

$$\begin{aligned} A_{h(^1S_0) \rightarrow P_1 P_2 P_3}(s_{12}, s_{13}, s_{23}) &= \Lambda_{12}(s, s_{12}) + \Lambda_{13}(s, s_{13}) + \Lambda_{23}(s, s_{23}) \\ &+ A_{12}(s, s_{12}) + A_{13}(s, s_{13}) + A_{23}(s, s_{23}), \end{aligned} \quad (89)$$

where the terms $A_{ij}(s, s_{ij})$ describe processes with interactions of all particles, P_1 , P_2 and P_3 . The term $A_{12}(s, s_{12})$ reads:

$$\begin{aligned} A_{12}(s, s_{12}) &= \left(B_{13-12}(s, s_{12}) + B_{23-12}(s, s_{12}) \right) \\ &\times \sum_n \frac{g_{12}^{(n)2}}{M_n^2 - s_{12}} f(s_{12}) \frac{1}{1 - b(s_{12}) \sum_{n'} \frac{g_{12}^{(n')2}}{M_{n'}^2 - s_{12}}}, \\ B_{i3-12}(s_{12}) &= \int_{(m_1+m_2)^2}^{\infty} \frac{ds'_{12}}{\pi} \langle \Lambda_{i3}(s, s'_{i3}) + A_{i3}(s, s'_{i3}) \rangle_{12}^{(0)} \frac{\rho_{12}^{(0)}(s'_{12}) f(s'_{12})}{s'_{12} - s_{12} - i0}. \end{aligned} \quad (90)$$




Figure 7: Complex- M plane for the $(IJ^{PC} = 00^{++})$ mesons. The dashed line encircles the part of the plane where the K -matrix analysis [4, 5, 6] reconstructs the analytical K -matrix amplitude: in this area the poles corresponding to resonances $f_0(980)$, $f_0(1300)$, $f_0(1500)$, $f_0(1750)$ and the broad state $f_0(1200 - 1600)$ are located. Beyond this area, in the low-mass region, the pole of the light σ -meson is located (shown by the point the position of pole, $M = (430 - i320)$ MeV, corresponds to the result of N/D analysis ; the crossed bars stand for σ -meson pole found in [9, 28]). In the high-mass region one has resonances $f_0(2030)$, $f_0(2100)$, $f_0(2340)$ [7]. Solid lines stand for the cuts related to the thresholds $\pi\pi$, $\pi\pi\pi\pi$, $K\bar{K}$, $\eta\eta$, $\eta\eta'$.

Analogous relations for $A_{13}(s, s_{13})$ and $A_{23}(s, s_{23})$ give us three non-homogenous equations for three amplitudes thus solving in principle the problem of construction of the three-body amplitude under the constraints of analyticity and unitarity.

The equations written here require a comment. We realise the convergence of the loop diagrams with the help of cutting vertices or, what is the same, the universal form factor. The convergence of a loop diagram can be realised in other ways as well. For example, in [19, 20] a special cutting function was introduced into the integrands; one may use the subtraction procedure as it is done in [23, 21]. The technical variations are of no importance, the only essential point is that for the convergence of the considered diagrams we have to introduce additional parameters – here it is the form factor slope μ^2 , see Eq. (84) and the corresponding discussion.

Miniconclusion

In this Appendix we have presented some characteristic features of the spectral integral equations for three-body systems. The technique can be used both for the determination of levels of compound systems and their wave functions (for instance, in the method of an accounting of the leading singularities [24] – this method was applied to the three-nucleon systems, H_3 and He_3 , and for determination of analytical properties of multiparticle production amplitudes when the produced resonances are studied.

We do not present here formulas for the reactions we have studied — the formulas are rather cumbersome. An example can be found, as it was mention above, in [20] where a set of equations for reactions $p\bar{p} \rightarrow \pi\pi\pi, \pi\eta\eta, \pi K\bar{K}$ was written.




Figure 8: Position of the f_2 -poles on the complex- M plane: states with dominant ${}^3P_2 n\bar{n}$ -component (full circle), ${}^3F_2 n\bar{n}$ -component (full triangle), ${}^3P_2 s\bar{s}$ -component (open circle), ${}^3F_2 s\bar{s}$ -component (open triangle); the position of the tensor glueball is shown by the open square. Mass regions studied by the groups L3 [29], PNPI-RAL [7, 30, 31, 32] and BNL [8, 33] are shown.

The presented technique may be especially convenient for the study of low-mass singularities in multiparticle production amplitudes. The long-lasting discussions on the sigma-meson observation, (see [25, 26, 27] and references therein) indicate that this is a problem of current interest.

5 Assignment of Mesons to Nonets

We present here the assignment of the mesons to $q\bar{q}$ nonets following [2].

In Figs. 7 and 8 we show the positions of the ($IJ^{PC} = 00^{++}$) and ($IJ^{PC} = 02^{++}$) resonances in the complex- M planes found in the previous analyses, see [2] for details.

In Eq. (91) we collected all considered meson $q\bar{q}$ states in nonets according to their $SU(3)_{flavour}$ attribution (the ${}^{2S+1}L_J$ states of $q\bar{q}$ mesons with $n = 1, 2, 3$ and 4). The sin-

glet and octet states, with the same values of the total angular momentum, are mixed.

$q\bar{q}$ -mesons	n=1				n=2			
	I=1	I=0	I=0	I= $\frac{1}{2}$	I=1	I=0	I=0	I= $\frac{1}{2}$
$^1S_0(0^{-+})$	$\pi(140)$	$\eta(547)$	$\eta'(958)$	$K(500)$	$\pi(1300)$	$\eta(1295)$	$\eta(1410)$	$K(1460)$
$^3S_1(1^{--})$	$\rho(775)$	$\omega(782)$	$\phi(1020)$	$K^*(890)$	$\rho(1460)$	$\omega(1430)$	$\phi(1650)$	
$^1P_1(1^{+-})$	$b_1(1229)$	$h_1(1170)$	$h_1(1440)$	$K_1(1270)$	$b_1(1620)$	$h_1(1595)$	$h_1(1790)$	$K_1(1650)$
$^3P_0(0^{++})$	$a_0(980)$	$f_0(980)$	$f_0(1300)$	$K_0(1425)$	$a_0(1474)$	$f_0(1500)$	$f_0(1750)$	$K_0(1820)$
$^3P_1(1^{++})$	$a_1(1230)$	$f_1(1282)$	$f_1(1426)$	$K_1(1400)$	$a_1(1640)$	$f_1(1518)$	$f_1(1780)$	
$^3P_2(2^{++})$	$a_2(1320)$	$f_2(1275)$	$f_2(1525)$	$K_2(1430)$	$a_2(1675)$	$f_2(1565)$	$f_2(1755)$	$K_2(1980)$
$^1D_2(2^{-+})$	$\pi_2(1676)$	$\eta_2(1645)$	$\eta_2(1850)$	$K_2(1800)$	$\pi_2(2005)$	$\eta_2(2030)$	$\eta_2(2150)$	
$^3D_1(1^{--})$	$\rho(1700)$	$\omega(1670)$		$K_1(1680)$	$\rho(1970)$	$\omega(1960)$		
$^3D_2(2^{--})$	$\rho_2(1940)$	$\omega_2(1975)$		$K_2(1580)$	$\rho_2(2240)$	$\omega_2(2195)$		$K_2(1773)$
$^3D_3(3^{--})$	$\rho_3(1690)$	$\omega_3(1667)$	$\phi_3(1854)$	$K_3(1780)$	$\rho_3(1980)$	$\omega_3(1945)$	$\phi_3(2140)$	
$^1F_3(3^{+-})$	$b_3(2032)$	$h_3(2025)$			$b_3(2245)$	$h_3(2275)$		
$^3F_2(2^{++})$	$a_2(2030)$	$f_2(2020)$	$f_2(2340)$		$a_2(2255)$	$f_2(2300)$	$f_2(2570)$	
$^3F_3(3^{++})$	$a_3(2030)$	$f_3(2050)$		$K_3(2320)$	$a_3(2275)$	$f_3(2303)$		
$^3F_4(4^{++})$	$a_4(2005)$	$f_4(2025)$	$f_4(2100)$	$K_4(2045)$	$a_4(2255)$	$f_4(2150)$	$f_4(2300)$	
$^1G_4(4^{-+})$	$\pi_4(2250)$	$\eta_4(2328)$		$K_4(2500)$				
$^3G_3(3^{--})$	$\rho_3(2240)$				$\rho_3(2510)$			
$^3G_4(4^{--})$								
$^3G_5(5^{--})$	$\rho_5(2300)$			$K_5(2380)$	$\rho_5(2570)$			
$^3H_6(6^{++})$	$a_6(2450)$	$f_6(2420)$						
$q\bar{q}$ -mesons	n=3				n=4			
	I=1	I=0	I=0	I= $\frac{1}{2}$	I=1	I=0	I=0	I= $\frac{1}{2}$
$^1S_0(0^{-+})$	$\pi(1800)$	$\eta(1760)$	$\eta(1880)$	$K(1830)$	$\pi(2070)$	$\eta(2010)$	$\eta(2190)$	
$^3S_1(1^{--})$	$\rho(1870)$	$\omega(\mathbf{1830})$	$\phi(1970)$		$\rho(2110)$	$\omega(2205)$	$\phi(2300)$	
$^1P_1(1^{+-})$	$b_1(1960)$	$h_1(1965)$	$h_1(2090)$		$b_1(2240)$	$h_1(2215)$		
$^3P_0(0^{++})$	$a_0(1780)$	$f_0(2040)$	$f_0(2105)$		$a_0(2025)$	$f_0(2210)$	$f_0(2340)$	
$^3P_1(1^{++})$	$a_1(1930)$	$f_1(1970)$	$f_1(2060)$		$a_1(2270)$	$f_1(2214)$	$f_1(2310)$	
$^3P_2(2^{++})$	$a_2(1950)$	$f_2(1920)$	$f_2(2120)$		$a_2(2175)$	$f_2(2240)$	$f_2(2410)$	
$^1D_2(2^{-+})$	$\pi_2(2245)$	$\eta_2(2248)$	$\eta_2(2380)$			$\eta_2(2520)$		
$^3D_1(1^{--})$	$\rho(2265)$	$\omega(2330)$						
$^3D_2(2^{--})$				$K_2(2250)$				
$^3D_3(3^{--})$	$\rho_3(2300)$	$\omega_3(2285)$	$\phi_3(2400)$					

(91)

In the lightest nonets we can determine mixing angles more or less reliably, but for the higher excitations the estimates of the mixing angles are very ambiguous. In addition, isoscalar states can contain significant glueball components. For these reasons, we give only the nonet ($9 = 1 \oplus 8$) classification of mesons. States that are predicted but not yet reliably established are shown in boldface.

Acknowledgments

We thank A.V. Anisovich, L.G. Dakhno, J. Nyiri, V.A. Nikonov, M.A. Matveev for helpful discussions. The paper was supported by the RFFI grant 07-02-01196-a.

References

- [1] V.V. Anisovich and A.V. Sarantsev *The analysis of reactions $\pi N \rightarrow two\ mesons + N$ within reggeon exchanges. 1.Fit and results*, in press, (2008).
- [2] A.V. Anisovich, V.V. Anisovich, J. Nyiri, V.A. Nikonov, M.A. Matveev and A.V. Sarantsev, *Mesons and Baryons. Systematization and Methods of Analysis*, World Scientific, Singapore, 2008.
- [3] V.V. Anisovich, M.N. Kobrinsky, J. Nyiri, and Yu.M. Shabelski, *Quark Model and High Energy Collisions*, 2nd edition, World Scientific, Singapore, 2004.
- [4] V.V. Anisovich and A.V. Sarantsev, Phys. Lett. B **382**, 429 (1996).
- [5] V.V. Anisovich, A.A. Kondashov, Yu.D. Prokoshkin, S.A. Sadovsky, and A.V. Sarantsev, Yad. Fiz. **60**, 1489 (2000) [Physics of Atomic Nuclei **60**, 1410 (2000)]; hep-ph/9711319.
- [6] V.V. Anisovich and A.V. Sarantsev, Eur. Phys. J. A **16**, 229 (2003).
- [7] A.V. Anisovich, C.A. Baker, C.J. Batty, *et al.*, Phys. Lett. B **449**, 114 (1999); B **452**, 173 (1999); B **452**, 180 (1999); B **452**, 187 (1999); B **472**, 168 (2000); B **476**, 15 (2000); B **477**, 19 (2000); B **491**, 40 (2000); B **491**, 47 (2000); B **496**, 145 (2000); B **507**, 23 (2001); B **508**, 6 (2001); B **513**, 281 (2001); B **517**, 261 (2001); B **517**, 273 (2001); Nucl. Phys. A **651**, 253 (1999); A **662**, 319 (2000); A **662**, 344 (2000).
- [8] S. J. Lindenbaum and R. S. Longacre, Phys. Lett. B **274**, 492 (1992).
- [9] V.V. Anisovich, UFN **174**, 49 (2004) [Physics-Uspekhi **47**, 45 (2004)].
- [10] A.V. Anisovich, V.V. Anisovich, V.N. Markov, M.A. Matveev, and A.V. Sarantsev, J. Phys. G: Nucl. Part. Phys. **28**, 15 (2002).
- [11] V.V. Anisovich, D.V. Bugg, A.V. Sarantsev, B.S. Zou, Talk at Symposium "*NN interaction (annihilation and scattering)*", October 1993, Moscow; Yad. Fiz. **57**, 1666 (1994) [Phys. Atom. Nucl. **57**, 1595 (1994)].
- [12] V.V. Anisovich, *et al.*, Phys. Lett. B **323**, 233 (1994); C. Amsler, *et al.*, Phys. Lett. B **342**, 433 (1995); B **355**, 425 (1995).
- [13] V.V. Anisovich, D.V. Bugg, A.V. Sarantsev, and B.S. Zou, Phys. Rev. D **50**, 1972 (1994).

- [14] V.V. Anisovich, D.S. Armstrong, I. Augustin, *et al.*, (Crystal Barrel Collab.), Phys. Lett. B **323**, 233 (1994).
- [15] C. Amsler, V. V. Anisovich, D.S. Armstrong, *et al.*, (Crystal Barrel Collab.), Phys. Lett. B **333**, 277 (1994).
- [16] G.V. Skornyyakov and K.A. Ter-Martirosyan, ZhETP **31**, 775 (1956);
G.S. Danilov, ZhETP **40**, 498 (1961); **42**, 1449 (1962).
- [17] L.D. Faddeev, ZhETP **41**, 1851 (1961).
- [18] V.V. Anisovich and L.G. Dakhno, Phys. Lett. **10**, 221 (1964).
- [19] A.V. Anisovich, Yad. Fiz. **58**, 1467 (1995) [Phys. Atom. Nucl. **58**, 1383 (1995)].
- [20] A.V. Anisovich, Yad. Fiz. **66**, 175 (2003) [Phys. Atom. Nucl. **66**, 172 (2003)].
- [21] V.V. Anisovich and A.A. Anselm, UFN **88**, 287 (1966) [Sov. Phys. Usp. **88**, 117 (1966)].
- [22] I.J.R. Aitchison and R. Pasquier, Phys. Rev. **152**, 1274 (1966).
- [23] V.V. Anisovich, A.A. Anselm, and V.N. Gribov, Nucl. Phys. **38**, 132 (1962).
- [24] A.V. Anisovich, V.V. Anisovich, Yad. Fiz. **53**, 1485 (1991) [Phys. Atom. Nucl. **53**, 915 (1991)]
- [25] M. Ablikim *et al.*, Phys. Lett. B **598**, 149 (2004).
- [26] D.V. Bugg, Phys. Rep. **397**, 257 (2004).
- [27] Z.Y. Zhou, *et al.*, JHEP:0502043 (2005).
- [28] V.V. Anisovich and V. A. Nikonov, Eur. Phys. J. A **8**, 401 (2000).
- [29] V.A. Schegelsky, A.V. Sarantsev, V.A. Nikonov, L3 Note 3001, October 27, 2004.
- [30] V.V. Anisovich, Pis'ma v ZhETF, **80**, 845 (2004) [JETP Letters, **80**, 715 (2004)].
- [31] V.V. Anisovich and A.V. Sarantsev, Pis'ma v ZhETF, **81**, 531 (2005), [JETP Letters, **81**, 417 (2005)].
- [32] V.V. Anisovich, M.A. Matveev, J. Nyiri, and A.V. Sarantsev, Int. J. Mod. Phys. A **20**, 6327 (2005); Yad. Fiz. **69**, 542 (2000) [Phys. Atom. Nuclei **69**, 520 (2006)].
- [33] R.S. Longacre and S.J. Lindenbaum, Report BNL-72371-2004.